

Clinical Policy: Histrelin Acetate (Vantas, Supprelin LA)

Reference Number: AZ.CP.PHAR.172

Effective Date: 12.01.21

Last Review Date: 05.23

Line of Business: Arizona Medicaid (AzCH-CCP and Care1st)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Histrelin acetate (Vantas[®] and Supprelin LA[®]) is a gonadotropin-releasing hormone (GnRH) agonist.

AHCCCS preferred drugs in this class include: Lupron Depot and Lupron Depot—Ped.

AHCCCS non-preferred drugs in this class include: Vantas, Supprelin LA.

FDA Approved Indication(s)

Vantas is indicated for the palliative treatment of advanced prostate cancer.

Supprelin LA is indicated for the treatment of children with central precocious puberty (CPP).

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of Arizona Complete Health-Complete Care Plan and Care1st that Vantas and Supprelin LA are **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Prostate Cancer (must meet all):

1. Diagnosis of prostate cancer;
2. Request is for Vantas, and medical justification supports inability to use Lupron Depot* (e.g., contraindications to the excipients);
**Prior authorization may be required for Lupron Depot.*
3. Prescribed by or in consultation with an oncologist or urologist;
4. Age \geq 18 years;
5. Request meets one of the following (a or b):*
 - a. Dose does not exceed 50 mg per 12 months (one implant per year);
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration: 12 months

B. Central Precocious Puberty (must meet all):

1. Diagnosis of CPP confirmed by all of the following (a, b, and c):
 - a. Elevated basal luteinizing hormone (LH) level > 0.2 - 0.3 mIU/L (dependent on type of assay used) and/or elevated leuprolide-stimulated LH level > 3.3 - 5 IU/L (dependent on type of assay used);
 - b. Difference between bone age and chronological age was > 1 year (bone age-chronological age);
 - c. Age at onset of secondary sex characteristics is < 8 years if female, or < 9 years if male;
2. Request is for Supprelin LA, and medical justification supports inability to use Lupron Depot-Ped (e.g., contraindications to the excipients Lupron Depot-Ped).
**Prior authorization may be required for Lupron Depot-Ped.*
3. Prescribed by or in consultation with a pediatric endocrinologist;
4. Member meets one of the following age requirements (a or b):
 - a. Female: 2 - 11 years;
 - b. Male: 2 - 12 years;
5. Dose does not exceed 50 mg per 12 months (one implant per year).

Approval duration: 12 months

C. Gender Dysphoria, Gender Transition (off-label) (must meet all):

1. Diagnosis of gender dysphoria or request is for gender transition;
2. Prescribed by or in consultation with an endocrinologist and a provider with expertise in gender dysphoria and transgender medicine based on a certified training program or affiliation with local transgender health services (e.g., mental health professional such as psychologist, psychiatrist, see *Appendix D*);
3. Request is for Supprelin LA and medical justification supports inability to use Lupron Depot-Ped (e.g., contraindications to the excipients Lupron Depot-Ped).
**Prior authorization may be required for Lupron Depot-Ped*
4. Age and pubertal development - meets (a or b):
 - a. Member is < 18 years of age and has reached or passed through Tanner Stage 2*;
**Age ranges approximating Tanner Stage 2 pubertal development extend from 8 to 13 years of age in girls and 9 to 14 years of age in boys.*
 - b. Member is ≥ 18 years of age and has failed to achieve physiologic hormone levels with gender-affirming hormonal therapy (e.g., estrogen, testosterone) unless contraindicated or clinically significant adverse effects are experienced;
5. Member demonstrates understanding of expected GnRH analogue treatment outcomes and has given consent for such treatment;
6. If member has a psychiatric comorbidity, member is followed by mental health provider;
7. Psychosocial support will be provided during treatment;
8. Dose is within FDA maximum limit for any FDA-approved indication (*see Section V*) or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

Approval duration: 12 months

D. Other diagnoses/indications

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): AZ.CP.PMN.53 for Arizona Medicaid.

II. Continued Therapy

A. Prostate Cancer (must meet all):

1. Currently receiving medication via Centene benefit, or documentation supports that member is currently receiving Vantas for prostate cancer and has received this medication for at least 30 days;
2. Request is for Vantas, and medical justification supports inability to use Lupron Depot* (e.g., contraindications to the excipients);
**Prior authorization may be required for Lupron Depot.*
3. Member is responding positively to therapy;
4. If request is for a dose increase, request meets one of the following (a or b):*
 - a. New dose does not exceed 50 mg per 12 months (one implant per year);
 - b. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration: 12 months

B. Central Precocious Puberty (must meet all):

1. Currently receiving medication via Centene benefit or member has previously met all initial approval criteria;
2. Request is for Supprelin LA;
3. Member is responding positively to therapy as evidenced by, including but not limited to, improvement in any of the following parameters: decreased growth velocity, cessation of menses, softening of breast tissue or testes, arrested pubertal progression;
4. Member meets one of the following age requirements (a or b):
 - a. Female: ≤ 11 years;
 - b. Male: ≤ 12 years;
5. If request is for a dose increase, new dose does not exceed 50 mg per 12 months (one implant per year).

Approval duration: 12 months

C. Gender Dysphoria, Gender Transition (off-label) (must meet all):

1. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
2. Member is responding positively to therapy;
3. If request is for a dose increase, new dose is within FDA maximum limit for any FDA-approved indication (*see Section V*) or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

Approval duration: 12 months

D. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.
Approval duration: Duration of request or 6 months (whichever is less); or
2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): AZ.CP.PMN.53 for Arizona Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – AZ.CP.PMN.53 for Arizona Medicaid.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

CPP: central precocious puberty	LH: luteinizing hormone
FDA: Food and Drug Administration	NCCN: National Comprehensive Cancer Network
GnRH: gonadotropin-releasing hormone	

Appendix B: Therapeutic Alternatives

Drug Name	Indication	Dosing Regimen	Maximum Dose
Leuprolide acetate injection Leuprolide acetate (Lupron Depot-Ped 7.5, 11.25, 15 [1 mo]; 11.25, 30 [3 mo])	CPP	Leuprolide acetate (SC): • Diagnostic: 20 mcg/kg or as needed; • Treatment: Initial: 50 mcg/kg/day; titrate dose upward by 10 mcg/kg/day if downregulation is not achieved (higher mg/kg doses may be required in younger children).	See regimen
		Lupron Depot-Ped (IM): Monthly administration weight-based starting dose: 7.5 mg (≤ 25 kg), 11.25 mg (> 25 to 37.5 kg), 15 mg (> 37.5 kg) (increase as needed to 15 mg/month); 3-month administration: 11.25 mg or 30 mg	See regimen

Therapeutic alternatives are listed as Brand name[®] (generic) when the drug is available by brand name only and generic (Brand name[®]) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): hypersensitivity to GnRH, GnRH agonist analogs; pregnancy
- Boxed warning(s): none reported

Appendix D: General Information

- World Professional Association for Transgender Health (WPATH) offers their Global Education Institute (GEI) Certified Training Courses: Best Practices in Transgender Medical and Mental Health Care. Additionally, the following link provides a search tool to locate WPATH member providers: <https://www.wpath.org/provider/search>
- Transgender Care Therapy Certification Training is also offered by the International Transgender Certification Association (ITCA). Professionals with expertise in transgender care can be located using the following search tool: <https://transgendercertification.com/locate-a-professional/>
- The draft of WPATH Standards of Care Version 8 are available and open for public comment. These standards of care recommend that adolescents are managed by a multidisciplinary care team that involves both medical and mental health professionals. The list of key disciplines includes but is not limited to: adolescent medicine/primary care, endocrinology, psychology, psychiatry, speech/language pathology, fertility, social work, support staff, and the surgical team. The need to include a healthcare professional with some expertise in mental health does not dictate the inclusion of a psychologist, psychiatrist or social work in every assessment. Instead, a general practitioner, nurse or other qualified clinician could fulfill this requirement as long as they have sufficient expertise to diagnose gender incongruence, recognize mental health concerns, distinguish between these concerns and gender dysphoria, incongruence or diversity, assist a transgender person in care planning and preparing for gender affirmative medical and surgical treatments, and refer to a mental health professional if needed.

V. Dosage and Administration

Drug Name	Indication	Dosing Regimen	Maximum Dose
Histrelin acetate (Supprelin LA)	CPP	1 implant (50 mg) SC for 12 months	1 implant per 12 months
Histrelin acetate (Vantas)	Prostate cancer - palliative therapy	1 implant (50 mg) SC for 12 months	1 implant per 12 months

VI. Product Availability

Drug Name	Availability
Histrelin acetate (Supprelin LA)	Implant: 50 mg (approximately 65 mcg histrelin acetate per day over 12 months)
Histrelin acetate (Vantas)	Implant: 50 mg (approximately 50 mcg histrelin acetate per day over 12 months)

VII. References

1. Vantas Prescribing Information. Malvern, PA: Endo Pharmaceuticals Solutions, Inc.; December 2020. Available at https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/021732s023lbl.pdf. Accessed July 28, 2022.
2. Supprelin LA Prescribing Information. Malvern, PA: Endo Pharmaceuticals Solutions, Inc.; April 2022. Available at www.supprelinla.com. Accessed July 28, 2022.
3. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at nccn.org. Accessed July 28, 2022.

4. National Comprehensive Cancer Network. Prostate Cancer Version 4.2022. Available at https://www.nccn.org/professionals/physician_gls/pdf/prostate.pdf. Accessed July 28, 2022.
 5. Kaplowitz P, Bloch C. Evaluation and referral of children with signs of early puberty. *Pediatrics*. 2016; 137(1): e20153732.
 6. Carel JC, Eugster EA, Rogol A, et al. Consensus statement on the use of gonadotropin-releasing hormone analogs in children. *Pediatrics*. 2009;123(4):e752. Epub 2009 Mar 30.
 7. Krishna KB, Fuqua JS, Rogol AD, et al. Use of gonadotropin-releasing hormone analogs in children: update by an International Consortium. *Horm Res Paediatr* 2019;91:357–372. DOI: 10.1159/000501336.
 8. Silverman LA, Neely EK, Kletter GB, et al. Long-term continuous suppression with once-yearly histrelin subcutaneous implants for the treatment of central precocious puberty: a final report of a phase 3 multicenter trial. *J Clin Endocrinol Metab*. 2015;100(6):2354-2363
- Gender Dysphoria*
9. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 5th ed. Arlington, VA: American Psychiatric Association Publishing; 2013.
 10. Hembree WC, Cohen-Kettenis PT, Gooren L, et al. Endocrine treatment of gender-dysphoric/gender-incongruent persons: An Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab*, November 2017, 102(11):3869–3903.
 11. Rafferty J. Ensuring comprehensive care and support for transgender and gender-diverse children and adolescents. Policy statement. American Academy of Pediatrics. *Pediatrics*; 142(4), October 2018:e20182162.
 12. Deutsch MB. Guidelines for the primary and gender-affirming care of transgender and gender nonbinary people. Center of Excellence for Transgender Health. Department of Family & Community Medicine, University of California, San Francisco; 2nd Ed, published June 17, 2016.
 13. Coleman E, Bockting W, Botzer M, et al. Standards of care for the health of transsexual, transgender, and gender nonconforming people. WPATH: World Professional Association for Transgender Health. 7th version; 2012. Available at https://www.wpath.org/media/cms/Documents/SOC%20v7/SOC%20V7_English2012.pdf?t=1613669341. Accessed July 21, 2022.
 14. Wylie KR, Fung R, Boshier C, Rotchell M. Recommendations of endocrine treatment for patients with gender dysphoria. *Sexual and Relationship Therapy* Vol. 24, No. 2, May 2009, 175–187.
 15. Emmanual M, Bokor BR. *Tanner Stages*. Treasure Island, FL: StatPearls Publishing; 2019 Jan. Available at <https://www.ncbi.nlm.nih.gov/books/NBK470280/>. Last update: May 13, 2019. Accessed July 21, 2022.
 16. Micromedex® Healthcare Series [Internet database]. Greenwood Village, Colo: Thomson Healthcare. Updated periodically. Accessed July 21, 2022.
 17. WPATH: World Professional Association for Transgender Health Standards of Care Version 8 Draft. Available at: <https://www.wpath.org/soc8>. Accessed July 21, 2022.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-

CLINICAL POLICY
Histrelin Acetate



date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J9225	Histrelin implant (Vantas), 50 mg
J9226	Histrelin implant (Supprelin LA) 50 mg
J1675	Injection, histrelin acetate, 10 micrograms

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created for AHCCCS preference of Lupron Depot for dx of CPP.	11.01.21	11.21
For CPP, clarified requirement for trial and failure of Lupron Depot-Ped from previously stated Lupron Depot; HCPCS code added for J1675 generic histrelin; references reviewed and updated.	02.07.23	02.23
Added off-label use criteria for gender dysphoria or gender transition; for Continued Therapy for Prostate Cancer, added requirement for trial and failure of Lupron Depot.	03.20.23	05.23

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan

CLINICAL POLICY

Histrelin Acetate



retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©YEAR 2021 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.